RULES OF THUMB IDENTIFIED FROM LITERATURE

SOWING	Seeding window is between Anzac Day and not beyond the June long weekend - NSW	Sow wheat to the depth of a matchbox on its side - SA	Only sow after ryegrass seed has germinated - SA	Fallow when the growing season gets less than 230 mm of rain, and soil has over 30 mm of stored water - WA	Fallow paddocks are lost potential - SA
FERTILISATION	Apply 4kg of phosphorous for every tonne of last year's crop - SA	Topdress if there is a greater than 5mm rain forecast - VIC	N fertiliser at 40-80kg per hectare (N/ha) has been found ideal for most growing condititions - WA	Nitrogen fertilizer loss is about 25% in low- medium rainfall, 32% in high rainfall, and 41% in irrigated - VIC	Where yields are consistently >2.5 t/ha, N applications can be delayed until stem elongation - SA
				Ů	J
SOIL HEALTH	Apply 2.5t/ha of lime in about every 10 years - VIC	Lime isn't economic unless pH is less than 4.5 or 4.8 and unless an acid sensitive plant like lucerne is grown - VIC	Apply lime once pH drops below 5 - VIC	Apply first remedial rate of lime approximately 150-200 kg lime/ha/year - VIC	Avoid cultivation if soil is too dry or too wet - VIC
		Ů			
CROP ROTATION	Don't follow one crop with another from the same family - ALL	Don't follow one heavy feeder with another heavy feeder - ALL	Do plant a green manure crop at least once in every 4 seasons to replenish your soil - ALL	Avoid planting certain groups of plants in the same spot year after year - ALL	Faba beans put the most N back into the paddock - VIC
MARKETING & FINANCIAL	Sell 1/3 grain at sowing, 1/3 later in the season and 1/3 at harvest - SA	\$300 is a good price for wheat - SA	Equity of 80% or more is comfortable - SA	You want to pay off new farms in 2 seasons - SA	If you can't pay off new machinery in 3 or 4 seasons, then you can't afford it - SA

